

STABILITY IN TERMS OF HYERS–ULAM OF NON–LINEAR VOLTERRA FREDHOLM INTEGRO–DELAY DYNAMIC SYSTEM ON TIME SCALES WITH FRACTIONAL INTEGRABLE IMPULSES

SYED OMAR SHAH AND ZUBAIR KHAN

ABSTRACT. This manuscript presents Hyers–Ulam stability and Hyers–Ulam–Rassias stability results of non–linear Volterra Fredholm integro–delay dynamic system on time scales with fractional integrable impulses. Picard fixed point theorem is used for obtaining existence and uniqueness of solutions. By means of abstract Grönwall lemma, Grönwall’s inequality on time scales, we establish Hyers–Ulam stability and Hyers–Ulam–Rassias stability results. There are some primary lemmas, inequalities and relevant assumptions that helps in our stability results

1. INTRODUCTION

In 1940, in a talk in front of the mathematics club at the university of Wisconsin, Ulam [34,35] presented a famed question related to the stability of homomorphisms: “*With which requirements does an additive mapping near an approximate additive mapping exists?*”.

This question was answered by Hyers [11] for the case when G_1 and G_2 are assumed to be Banach spaces by using direct method. Since then, this interesting stability, initiated by Ulam and Hyers, is called Hyers–Ulam stability. In 1978, Rassias [21] extended Hyers–Ulam stability concept by introducing new function variables, so this kind of stability is known as the Hyers–Ulam–Rassias stability. In fact, the most interesting result was of Rassias [21] that weakens the condition for the bound of the norm of Cauchy difference $f(x + y) - f(x) - f(y)$. For further details and discussions, we recommend the book by Jung [13].

At the end of 19th century, a large number of researchers contributed to the stability idea of Ulam’s type for various types of differential equations. There are many advantages of Ulam’s type stability in tackling problems related to optimization techniques, numerical analysis, control theory and many more, in such situations to get an exact solution is challenging. For more details on Hyers–Ulam stability, see [12, 14, 18, 19, 22–26, 28, 29, 31, 38, 40, 42, 43, 45–47, 49].

There are several implications for simple differential equations. Anyhow, the circumstances rather change when a real world process undergoes with unexpected variations, like significant mechanical processes, blood flows, heart beats, changes in population, radio physics, pharmacokinetics, mathematical economy, chemical technology, electrical technology, chemistry, different engineering fields, control theory

2020 *Mathematics Subject Classification.* 34N05, 34D20, 34A37, 45M10, 45J05.

Key words and phrases. Hyers–Ulam stability, time scale, impulses, delay dynamic system, Grönwall’s inequality, abstract Grönwall lemma, Banach fixed point theorem.

and so on, see [4, 5, 17]. Such circumstances generate a differential equation, which is known as impulsive differential equation. More precisely, there are three parts of differential equations with impulse impact: an instantaneous impulsive differential equation, in which the impulse action is defined at certain discrete points; non-instantaneous impulsive differential equation, it establishes the effect of impulse on an interval; and the third one is an impulse rule, in which we define a distinct and well defined collection of impulse events having an active impulse equation.

Fractional differential and integral equations play a key role not only in mathematics but also in the modeling of various physical phenomena in physics, control systems and dynamical systems. In fact, fractional order derivatives and integrals are assumed to be more realistic and practical than derivatives and integrals of integral order. These are excellent tools to model genetic transformation and memory retention qualities of several systems and products.

It is to be noted that, the pioneers of the Ulam's type stability for impulsive ordinary differentiable equations are Wang *et al.* [36]. Following their own work, in 2014, they proved the Hyers–Ulam–Rassias stability and generalized Hyers–Ulam–Rassias stability of impulsive evolution equations on a compact interval [37] which then they extended for infinite impulses in the same paper. Wang and Zhang [39], initially studied nonlinear differential equations having fractional integrable impulses, which are more interesting. They presented four Bielecki–Ulam's type stabilities for this class of differential equations. Also Lin *et al.* [15] discussed the existence and stability results for impulsive integro-differential equations. The work of Wang *et al.* [39] was extended by Zada *et al.* [43] in which they discussed Hyers–Ulam stability of higher-order nonlinear differential equations with fractional integrable impulses. They established Bielecki–Ulam–Hyers–Rassias stability, generalized Bielecki–Ulam–Hyers–Rassias stability and Bielecki–Ulam–Hyers stability for this class of differential equations on a compact interval. Recently, Zada *et al.* [48] obtained very interesting results about the Hyers–Ulam stability of nonlinear impulsive Volterra integro-delay dynamic system on time scales.

However, despite the situations where only impulsive factor is involved or delay effects happened, we have a wide variety of evolutionary processes with both delay and impulsive effects. To model such phenomena which are subject to impulsive perturbations as the time delays, an impulsive delay differential equation is used.

The theory of dynamic equations on time scales has been rising fast and has acknowledged a lot of interest in recent years. This theory was introduced by Hilger [10] in 1988, with the inspiration to provide a unification of continuous and discrete calculus. For more details on time scales, see [1–3, 6–9, 16, 20, 28, 30–33, 41, 44, 47].

As far as we know, not too many results of stability of delay dynamic equations with impulses are analyzed by researchers. Although, to the extent of our knowledge, the stability observations of Ulam's type of non-linear Volterra Fredholm integro-delay dynamic system having integral impulsions of fractional order are not yet investigated.

Motivated by the work done in [30, 48], the utmost purpose of this manuscript is to find different Hyers–Ulam and Hyers–Ulam–Rassias outcomes of stability for

the following non-linear Volterra Fredholm integro–delay dynamic system with integrable impulses having fractional order

$$(1.1) \quad \begin{cases} \omega^\Delta(t) = M(t)\omega(t) + \int_{t_0}^t \mathcal{K}(t, s, \omega(s), \omega(h(s)))\Delta s \\ \quad + \int_a^b \mathcal{K}(t, s, \omega(s), \omega(h(s)))\Delta s, \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \\ \omega(t) = I_{t_i, t}^\alpha g_i(t, \omega(t), \omega(h(t))), \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \\ \omega(t) = \alpha(t), \quad t \in [s_0 - \lambda, s_0] \cap T_S, \\ \omega(t_0) = \alpha(t_0) = \omega_0, \end{cases}$$

where $\lambda > 0$, T_S is a time scale, $M(t)$ is a piecewise continuous regressive square matrix, $0 = t_0 = s_0 < t_1 \leq s_1 \leq t_2 < \dots < t_m \leq s_m \leq t_{m+1} = t_f$ are pre-fixed numbers, $\mathcal{K}(t, s, \omega(s), \omega(h(s)))$ is piecewise continuous operator on $\Gamma = \{(t, s, \omega) : t_0 \leq s \leq t \leq t_f, \omega \in \mathbb{R}^m\}$, $g_i : (t_i, s_i] \cap T_S \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$, $i = 1, 2, \dots, m$ are continuous functions, $\phi : [s_0 - \lambda, s_0] \cap T_S \rightarrow \mathbb{R}^n$ is history function and $I_{t_i, t}^\alpha g_i$ are the so called Riemann–Liouville integrals having fractional order $\alpha \in (0, 1)$, with the representation:

$$I_{t_i, t}^\alpha g_i(t, \omega(t), \omega(h(t))) = \frac{1}{\Gamma(\alpha)} \int_{t_i}^t (t - s)^{\alpha-1} g_i(s, \omega(s), \omega(h(s)))\Delta s.$$

Moreover, $(s_i, t_{i+1}] \cap T_S$, $(t_i, s_i] \cap T_S$, $[s_0 - \lambda, s_0] \cap T_S$ are non-empty sets and $h : [s_0 - \lambda, t_f] \cap T_S \rightarrow (s_i, t_{i+1}] \cap T_S$ is a delay function with the consumption of continuity, additionally $h(t) \leq t$.

2. PRELIMINARIES

In this section, we recall the main definitions and some basic notations of time scales calculus.

An arbitrary non-empty closed subset of real numbers T_S is called a time scale. The forward jump operator $\Theta : T_S \rightarrow T_S$, backward jump operator $\rho : T_S \rightarrow T_S$ and graininess operator $\mu : T_S \rightarrow [0, \infty)$, are defined by:

$$\Theta(s) = \inf\{t \in T_S : t > s\}, \quad \rho(s) = \sup\{t \in T_S : t < s\}, \quad \mu(s) = \Theta(s) - s,$$

respectively. An arbitrary $t \in T_S$ is called left scattered (resp. left dense) when $t < \rho(t)$ (resp. $t = \rho(t)$). While, in case of $t < \Theta(t)$ (resp. $\Theta(t) = t$), we call t right scattered (resp. right dense). For a time scale T_S , the set of all limiting points T_S^z is called the derived set and illustrated as follows:

$$T_S^z = \begin{cases} T_S \setminus (\rho(\sup T_S), \sup T_S], & \text{if } \sup T_S < \infty, \\ T_S, & \text{if } \sup T_S = \infty. \end{cases}$$

The function $\mathcal{W} : T_S \rightarrow \mathbb{R}$ is called right-dense continuous if it is continuous at every right dense point on T_S and its left sided limit exists at every left dense point on T_S . The function $\mathcal{W} : T_S \rightarrow \mathbb{R}$ is called regressive (resp. positively regressive) if $1 + \mu(t)\mathcal{W}(t) \neq 0$, (resp. $1 + \mu(t)\mathcal{W}(t) > 0$) $\forall t \in T_S^z$. The set of all right-dense continuous regressive functions (resp. right-dense continuous positively regressive

functions) will be denoted by $\mathcal{R}_{\mathcal{G}}(T_S)$ (resp. $\mathcal{R}_{\mathcal{G}}(T_S)^+$). The delta derivative of the function $W : T_S \rightarrow \mathbb{R}$ on $t \in T_S^z$, is given by

$$W^\Delta(t) = \lim_{s \rightarrow t, s \neq \Theta(t)} \frac{W(\Theta(t)) - W(s)}{\Theta(t) - s}.$$

For a rd-continuous function $W : T_S \rightarrow \mathbb{R}$, the Δ -integral is defined to be

$$\int_a^b W(t) \Delta t = w(b) - w(a), \text{ for all } a, b \in T_S,$$

where w is the anti-derivative of W , i.e., $w^\Delta = W$ on T_S^z .

For $p \in \mathcal{R}_{\mathcal{G}}(T_S)$, the generalized exponential function is defined by

$$e_p(a, b) = \exp \left(\int_a^b \alpha_{\mu(s)} p(s) \Delta s \right) \text{ for all } a, b \in T_S,$$

while,

$$\alpha_{\mu(t)} p(t) = \begin{cases} \frac{\text{Log}(1 + \mu(t)p(t))}{\mu(t)}, & \text{if } \mu(t) \neq 0, \\ p(t), & \text{if } \mu(t) = 0, \end{cases}$$

is the cylindrical transformation.

The fundamental matrix $\Psi_M(t, t_0)$ is the unique solution of the dynamic equation $\omega^\Delta(t) = M(t)\omega(t)$, $\omega(t_0) = \omega_0$, $t \in T_S^0$.

3. BASIC CONCEPTS AND REMARKS

Let $C(J, \mathbb{R}^n)$ (resp. $PC(J, \mathbb{R}^n)$) be the Banach space of all continuous functions (resp. the Banach space of piecewise continuous functions) with the norm $\|\omega\|_\infty = \sup_{t \in J} \|\omega(t)\|$, $J = [s_0 - \lambda, t_f] \cap T_S$ and \mathbb{R} represents the set of real numbers. Finally, we denote by $PC^1(J, \mathbb{R}^n) = \{\omega \in PC(J, \mathbb{R}^n) : \omega^\Delta \in PC(J, \mathbb{R}^n)\}$, the Banach space with norm $\|\omega\|_1 = \max\{\|\omega\|_\infty, \|\omega^\Delta\|_\infty\}$. Here, as usual we denote by $\|x\| = \sum_{i=1}^n |x_i|$ for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$. Consider the following inequalities,

$$(3.1) \quad \begin{cases} \left| \left| \phi^\Delta(t) - M(t)\phi(t) - \int_{t_0}^t \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s \right. \right. \\ \left. \left. - \int_a^b \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s \right| \leq \epsilon, \quad t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \right. \\ \left. \left| \left| \phi(t) - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right| \leq \epsilon, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \right. \end{cases}$$

$$(3.2) \quad \begin{cases} \left| \left| \phi^\Delta(t) - M(t)\phi(t) - \int_{t_0}^t \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s \right. \right. \\ \left. \left. - \int_a^b \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s \right| \leq \varphi(t), \quad t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \right. \\ \left. \left| \left| \phi(t) - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right| \leq \kappa, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \right. \end{cases}$$

where $\epsilon > 0$, $\kappa \geq 0$ and $\varphi \in PC(J, \mathbb{R}^+)$ is an increasing function.

Definition 3.1. Eq. (1.1) is said to be stable in the sense of Hyers–Ulam, if for every $\epsilon > 0$ there exists a positive number K such that for every $\phi \in PC^1(J, \mathbb{R}^n)$ satisfying (3.1), there exists a solution $\phi_0 \in PC^1(J, \mathbb{R}^n)$ of (1.1) such that $\|\phi_0(t) - \phi(t)\| \leq K\epsilon$ for all $t \in J$. Here K is a positive number that depends on ϵ and do not depend on f_i .

Definition 3.2. Eq. (1.1) is said to be stable in the sense of Hyers–Ulam–Rassias, provided for all $(\varphi, \kappa) \in PC(J, \mathbb{R}^+) \times \mathbb{R}^+$ there exists $M > 0$ such that for all $\phi \in PC^1(J, \mathbb{R}^n)$ satisfying (3.2), there exists a solution $\phi_0 \in PC^1(J, \mathbb{R}^n)$ of (1.1) such that the inequality $\|\phi_0(t) - \phi(t)\| \leq M\varphi(t)$ is true for all $t \in J$. Here $M > 0$ depends on (φ, κ) .

Definition 3.3. In a metric space $(X; d)$, a mapping $\Lambda : X \rightarrow X$ is said to be Picard operator if it has precisely a unique fixed point $x^* \in X$, so that for every $x \in X$, the sequence $\{\Lambda^{(n)}(x)\}_{n \in \mathbb{N}}$ converges to x^* .

Lemma 3.4 ([16]). *Suppose $\tau \in T_S^+$, $y, b \in \mathcal{R}_g(T_S^+)$, $p \in \mathcal{R}_g(T_S^+)^+$ and $c, b_k \in \mathbb{R}^+$, $k = 1, 2, \dots$, so*

$$y(t) \leq c + \int_{\tau}^t p(s)y(s)\Delta s + \sum_{\tau < t_k < t} b_k y(t_k),$$

implies

$$y(t) \leq c \prod_{\tau < t_k < t} (1 + b_k) e_p(t, \tau), \quad t \geq \tau.$$

Lemma 3.5 (Abstract Grönwall Lemma [27]). *Let (X, d, \leq) be an ordered metric space and let x^* be a fixed point for the increasing mapping $\Lambda : X \rightarrow X$. So, being arbitrary $x \in X$, $x \leq \Lambda(x)$ entails $x \leq x^*$ and $x \geq \Lambda(x)$ entails $x \geq x^*$, where x^* denotes the fixed point in Λ .*

Remark 3.6. A function $\phi \in PC^1(J, \mathbb{R}^n)$ satisfies inequality (3.1) (resp. inequality (3.2)) if and only if there exist a function $f \in PC^1(J, \mathbb{R}^n)$ and a finite sequence $\{f_k : k = 1, \dots, m\} \subset \mathbb{R}^n$ (dependent on ϕ) such that $\|f(t)\| \leq \epsilon$ for all $t \in J$ and $\|f_i\| \leq \epsilon$ (resp. $\|f_i\| \leq \kappa$) for every $i = 1, 2, \dots, m$ and

$$\left\{ \begin{array}{l} \phi^\Delta(t) = M(t)\phi(t) + \int_{t_0}^t \mathcal{K}(t, s, \phi(s), \phi(h(s)))\Delta s \\ + \int_a^b \mathcal{K}(t, s, \phi(s), \phi(h(s)))\Delta s + f(t), \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \\ \phi(t) = I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) + f_i, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m. \end{array} \right.$$

Lemma 3.7. *If $\phi \in PC^1(J, \mathbb{R}^n)$ satisfies inequality (3.1) (resp. inequality (3.2)), then the following inequalities*

$$\left\{ \begin{array}{l} \left\| \phi(t) - \phi_0 - \Psi_M(t, t_0)\phi_0 - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_{s_0}^s \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u \Delta s \right. \\ \left. - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_a^b \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u \Delta s \right. \\ \left. - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right\| \leq (Ct_f - Cs_i + m)\epsilon, \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 1, 2, \dots, m, \\ \left\| \phi(t) - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right\| \leq m\epsilon, \quad (\text{resp. } m\kappa), \\ t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \end{array} \right.$$

are true. Here C is the bound of fundamental matrix $\Psi_M(t, \Theta(s))$.

Proof. If $\phi \in PC^1(J, \mathbb{R}^n)$ satisfies (3.1), then by Remark 3.6, we have

$$(3.3) \quad \left\{ \begin{array}{l} \phi^\Delta(t) = M(t)\phi(t) + \int_{t_0}^t \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s \\ + \int_a^b \mathcal{K}(t, s, \phi(s), \phi(h(s))) \Delta s + f(t), \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \\ \phi(t) = I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) + f_i, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m. \end{array} \right.$$

Clearly the solution of (3.3) is given as

$$\phi(t) = \left\{ \begin{array}{l} \phi_0 + \Psi_M(t, t_0)\phi_0 + \int_{s_i}^t \Psi_M(t, \Theta(s)) \left(\int_{s_0}^s \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u \right. \\ \left. + \int_a^b \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u + f(s) \right) \Delta s + I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))), \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 1, 2, \dots, m, \\ I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) + f_i, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m. \end{array} \right.$$

For $t \in (s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$, we get

$$\begin{aligned} & \left\| \phi(t) - \phi_0 - \Psi_M(t, t_0)\phi_0 - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_{s_0}^s \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u \Delta s \right. \\ & \quad \left. - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_a^b \mathcal{K}(s, u, \phi(u), \phi(h(u))) \Delta u \Delta s - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right\| \\ & \leq \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \|f(s)\| ds + \sum_{i=1}^m \|f_i\| \\ & \leq (Ct - Cs_i + m)\epsilon \\ & \leq (Ct_f - Cs_i + m)\epsilon. \end{aligned}$$

Proceeding as above we derive

$$\left\| \phi(t) - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t))) \right\| \leq m\epsilon, \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m.$$

We have similar processions for (3.2). \square

4. MAIN RESULTS

Onward we will state our major results. The first solution to be establish is Hyers–Ulam stability. First we assume the following conditions:

(A₁) The function \mathcal{K} is piecewise continuous with the Lipschitz condition $\|\mathcal{K}(t, s, x_1, x_2) - \mathcal{K}(t, s, y_1, y_2)\| \leq \sum_{k=1}^2 L \|x_k - y_k\|$, $L > 0$, for all $t \in (s_i, t_{i+1}] \cap T_S$, $i = 0, 1, \dots, m$ and $x_k, y_k \in \mathbb{R}^n$, $k \in \{1, 2\}$;

(A₂) $g_i : (t_i, s_i] \cap T_S \times \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ satisfies the Lipschitz condition $\|g_i(t, u_1, u_2) - g_i(t, v_1, v_2)\| \leq \sum_{k=1}^2 L_{g_i} \|u_k - v_k\|$, $L_{g_i} > 0$, for all $t \in (t_i, s_i] \cap T_S$, $i = 1, 2, \dots, m$ and $u_1, u_2, v_1, v_2 \in \mathbb{R}^n$;

(A₃) $\left(\frac{2L_{g_i}}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \Delta s + 2CL \int_{s_i}^t \int_{s_0}^s \Delta u \Delta s + 2CL(t_f - s_i)(b - a) \right) < 1$, $i = 1, 2, \dots, m$;

(A₄) $\varphi \in PC(J, \mathbb{R}^+)$ is increasing so that for some $\rho > 0$,

$$\int_{t_0}^t \varphi(r) \Delta r \leq \rho \varphi(t).$$

Theorem 4.1. *If conditions (A₁) – (A₃) hold, then Eq. (1.1) has precisely a unique solution in $PC^1(J, \mathbb{R}^n)$.*

Proof. Determine an operator $\Lambda : PC(J, \mathbb{R}^n) \rightarrow PC(J, \mathbb{R}^n)$ as

$$(4.1) \quad (\Lambda\omega)(t) = \begin{cases} \alpha(t), & t \in [s_0 - \lambda, s_0] \cap T_S, \\ I_{t_i, s_i}^\alpha g_i(t, \omega(t), \omega(h(t))), & t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \quad \alpha \in (0, 1), \\ \alpha(t_0) + \Psi_M(t, t_0)\omega_0 + I_{t_i, s_i}^\alpha g_i(s_i, \omega(s_i), \omega(h(s_i))) \\ + \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_{s_0}^s \mathcal{K}(s, u, \omega(u), \omega(h(u))) \Delta u \Delta s \\ + \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_a^b \mathcal{K}(s, u, \omega(u), \omega(h(u))) \Delta u \Delta s, \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 1, 2, \dots, m, \quad \alpha \in (0, 1). \end{cases}$$

For any $\omega_1, \omega_2 \in PC(J, \mathbb{R}^n)$, $t \in (s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$, we have

$$\begin{aligned} & \|(\Lambda\omega_1)(t) - (\Lambda\omega_2)(t)\| \\ & \leq \|I_{t_i, s_i}^\alpha g_i(s_i, \omega_1(s_i), \omega_1(h(s_i))) - I_{t_i, s_i}^\alpha g_i(s_i, \omega_2(s_i), \omega_2(h(s_i)))\| \\ & \quad + \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_{s_0}^s \|\mathcal{K}(s, u, \omega_1(u), \omega_1(h(u)))\| \end{aligned}$$

$$\begin{aligned}
& - \mathcal{K}(s, u, \omega_2(u), \omega_2(h(u))) \Big\| \Delta u \Delta s \\
& + \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_a^b \left\| \mathcal{K}(s, u, \omega_1(u), \omega_1(h(u))) \right. \\
& \left. - \mathcal{K}(s, u, \omega_2(u), \omega_2(h(u))) \right\| \Delta u \Delta s \\
\leq & \frac{1}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|g_i(s, \omega_1(s), \omega_1(h(s))) \\
& - g_i(s, \omega_2(s), \omega_2(h(s)))\| \Delta s \\
& + L \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_{s_0}^s \|\omega_1(u) - \omega_2(u)\| \Delta u \Delta s \\
& + L \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_{s_0}^s \|\omega_1(h(u)) - \omega_2(h(u))\| \Delta u \Delta s \\
& + L \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_a^b \|\omega_1(u) - \omega_2(u)\| \Delta u \Delta s \\
& + L \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_a^b \|\omega_1(h(u)) - \omega_2(h(u))\| \Delta u \Delta s \\
\leq & \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|\omega_1(s) - \omega_2(s)\| \Delta s \\
& + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|\omega_1(h(s)) - \omega_2(h(s))\| \Delta s \\
& + 2CL \int_{s_i}^t \int_{s_0}^s \sup_{s_i \leq s \leq t_{i+1}} \|\omega_1(u) - \omega_2(u)\| \Delta u \Delta s \\
& + 2CL \int_{s_i}^t \int_a^b \sup_{s_i \leq s \leq t_{i+1}} \|\omega_1(u) - \omega_2(u)\| \Delta u \Delta s \\
\leq & \frac{2Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \sup_{t_i \leq s \leq s_i} \|\omega_1(s) - \omega_2(s)\| \Delta s \\
& + 2CL \int_{s_i}^t \int_{s_0}^s \sup_{s_i \leq s \leq t_{i+1}} \|\omega_1(s) - \omega_2(s)\| \Delta u \Delta s \\
& + 2CL \int_{s_i}^t \int_a^b \sup_{s_i \leq s \leq t_{i+1}} \|\omega_1(s) - \omega_2(s)\| \Delta u \Delta s \\
\leq & \left(\frac{2Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \Delta s + 2CL \int_{s_i}^t \int_{s_0}^s \Delta u \right. \\
& \left. + 2CL(t_f - s_i)(b - a) \right) \|\omega_1 - \omega_2\|_\infty.
\end{aligned}$$

According to **(A₃)**, we are dealing here with the strictly contractive operator on $(s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$, and hence a Picard operator on $PC(J, \mathbb{R}^n)$. Regarding to (4.1), it shows that the unique solution of Eq. (1.1) in $PC^1(J, \mathbb{R}^n)$ is in fact the

unique fixed point of this operator. □

Theorem 4.2. *If conditions (\mathbf{A}_1) – (\mathbf{A}_3) hold, then Eq. (1.1) has Hyers–Ulam stability on J .*

Proof. Assume that (3.1) has a solution $\phi \in PC^1(J, \mathbb{R}^n)$. Then for dynamic equation

$$\begin{cases} \omega^\Delta(t) = M(t)\omega(t) + \int_{t_0}^t \mathcal{K}(t, s, \omega(s), \omega(h(s)))\Delta s \\ + \int_a^b \mathcal{K}(t, s, \omega(s), \omega(h(s)))\Delta s, \quad t \in (s_i, t_{i+1}] \cap T_S, \quad i = 0, 1, \dots, m, \\ \omega(t) = I_{t_i, t}^\alpha g_i(t, \omega(t), \omega(h(t))), \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \\ \omega(t) = \phi(t), \quad t \in [s_0 - \lambda, s_0] \cap T_S, \\ \omega(t_0) = \phi(t_0) = \omega_0, \end{cases}$$

we have the unique solution

$$\omega(t) = \begin{cases} \phi(t), \quad t \in [s_0 - \lambda, s_0] \cap T_S, \\ I_{t_i, s_i}^\alpha g_i(t, \omega(t), \omega(h(t))), \quad t \in (t_i, s_i] \cap T_S, \quad i = 1, 2, \dots, m, \quad \alpha \in (0, 1), \\ \phi(t_0) + \Psi_M(t, t_0)\omega_0 + I_{t_i, s_i}^\alpha g_i(s_i, \omega(s_i), \omega(h(s_i))) \\ + \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_{s_0}^s \mathcal{K}(s, u, \omega(u), \omega(h(u)))\Delta u \Delta s \\ + \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_a^b \mathcal{K}(s, u, \omega(u), \omega(h(u)))\Delta u \Delta s, \\ t \in (s_i, t_{i+1}] \cap T_S, \quad i = 1, 2, \dots, m. \end{cases}$$

We observe that for all $t \in (s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$, using Lemma 3.7, we have

$$\begin{aligned} & \|\phi(t) - \omega(t)\| \\ & \leq \|\phi(t) - \phi_0 - \Psi_M(t, t_0)\phi_0 - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_{s_0}^s \mathcal{K}(s, u, \phi(u), \phi(h(u)))\Delta u \\ & \quad - \int_{s_i}^t \Psi_M(t, \Theta(s)) \int_a^b \mathcal{K}(s, u, \phi(u), \phi(h(u)))\Delta u \Delta s \\ & \quad - I_{t_i, t}^\alpha g_i(t, \phi(t), \phi(h(t)))\| + \|I_{t_i, s_i}^\alpha g_i(s_i, \phi_1(s_i), \phi_1(h(s_i))) \\ & \quad - I_{t_i, s_i}^\alpha g_i(s_i, \phi_2(s_i), \phi_2(h(s_i)))\| \\ & \quad + \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_{s_0}^s \|\mathcal{K}(s, u, \phi_1(u), \phi_1(h(u))) \\ & \quad - \mathcal{K}(s, u, \phi_2(u), \phi_2(h(u)))\| \Delta u \Delta s \\ & \quad + \int_{s_i}^t \|\Psi_M(t, \Theta(s))\| \int_a^b \|\mathcal{K}(s, u, \phi_1(u), \phi_1(h(u))) \\ & \quad - \mathcal{K}(s, u, \phi_2(u), \phi_2(h(u)))\| \Delta u \Delta s \end{aligned}$$

$$\begin{aligned}
&\leq (m + Ct_f - Cs_i)\epsilon + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|\phi_1(s) - \phi_2(s)\| \Delta s \\
&\quad + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|\phi_1(h(s)) - \phi_2(h(s))\| \Delta s \\
&\quad + CL \int_{s_i}^t \int_{s_0}^s \|\phi_1(u) - \phi_2(u)\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_{s_0}^s \|\phi_1(h(u)) - \phi_2(h(u))\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_a^b \|\phi_1(u) - \phi_2(u)\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_a^b \|\phi_1(h(u)) - \phi_2(h(u))\| \Delta u \Delta s.
\end{aligned}$$

Next, we show that the operator $T : PC(J, \mathbb{R}^+) \rightarrow PC(J, \mathbb{R}^+)$ given below is an increasing Picard operator:

$$\begin{aligned}
(Tg)(t) &= (m + Ct_f - Cs_i)\epsilon + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g(s) \Delta s \\
&\quad + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g(h(s)) \Delta s + CL \int_{s_i}^t \int_{s_0}^s g(u) \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_{s_0}^s g(h(u)) \Delta u \Delta s + CL \int_{s_i}^t \int_a^b g(u) \Delta u \Delta s + CL \int_{s_i}^t \int_a^b g(h(u)) \Delta u \Delta s.
\end{aligned}$$

For any $g_1, g_2 \in PC(J, \mathbb{R}^+)$, $t \in (s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$, we have

$$\begin{aligned}
&\|(Tg_1)(t) - (Tg_2)(t)\| \leq \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|g_1(s) - g_2(s)\| \Delta s \\
&\quad + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \|g_1(h(s)) - g_2(h(s))\| \Delta s \\
&\quad + CL \int_{s_i}^t \int_{s_0}^s \|g_1(u) - g_2(u)\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_{s_0}^s \|g_1(h(u)) - g_2(h(u))\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_a^b \|g_1(u) - g_2(u)\| \Delta u \Delta s \\
&\quad + CL \int_{s_i}^t \int_a^b \|g_1(h(u)) - g_2(h(u))\| \Delta u \Delta s \\
&\leq \frac{2Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \sup_{t_i \leq s \leq s_i} \|g_1(s) - g_2(s)\| \Delta s \\
&\quad + 2CL \int_{s_i}^t \int_{s_0}^s \sup_{s_i \leq s \leq t_{i+1}} \|g_1(s) - g_2(s)\| \Delta u \Delta s
\end{aligned}$$

$$\begin{aligned}
& +2CL \int_{s_i}^t \int_a^b \sup_{s_i \leq s \leq t_{i+1}} \|g_1(s) - g_2(s)\| \Delta u \Delta s \\
\leq & \left(\frac{2Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \Delta s + 2CL \int_{s_i}^t \int_{s_0}^s \Delta u \Delta s \right. \\
& \left. + 2CL(t_f - s_i)(b - a) \right) \|g_1 - g_2\|_\infty.
\end{aligned}$$

Again according to **(A₃)**, we are dealing here with the strictly contractive operator on $(s_i, t_{i+1}] \cap T_S$, $i = 1, 2, \dots, m$ and hence a Picard operator on $PC(J, \mathbb{R}^+)$. Banach fixed point theorem imply, T is Picard operator having unique fixed point $g^* \in PC(J, \mathbb{R}^+)$ i.e.,

$$\begin{aligned}
g^*(t) &= (m + Ct_f - Cs_i)\epsilon + \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g^*(s) \Delta s \\
&+ \frac{Lg_i}{\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g^*(h(s)) \Delta s + CL \int_{s_i}^t \int_{s_0}^s g^*(u) \Delta u \Delta s \\
&+ CL \int_{s_i}^t \int_{s_0}^s g^*(h(u)) \Delta u \Delta s + CL \int_{s_i}^t \int_a^b g^*(u) \Delta u \Delta s \\
&+ CL \int_{s_i}^t \int_a^b g^*(h(u)) \Delta u \Delta s.
\end{aligned}$$

As, g^* is increasing, therefore $g^*(h(t)) \leq g^*(t)$, $(m + Ct_f - Cs_i) \leq \delta$ for some $\delta > 0$, $i = 1, 2, \dots, m$ and $\int_a^b g^*(h(u)) \Delta u = \varrho\epsilon$ be a fixed value, then we can write

$$\begin{aligned}
g^*(t) &\leq \delta\epsilon + \frac{2Lg_i}{m\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g^*(s) \Delta s + 2CL \int_{s_0}^t \int_{s_0}^s g^*(u) \Delta u \Delta s \\
&+ 2CL \int_{s_0}^t \varrho\epsilon \Delta s \\
&\leq \delta\epsilon + \frac{2Lg_i}{m\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} g^*(s) \Delta s + 2CL \int_{s_0}^t \int_{s_0}^s g^*(u) \Delta u \Delta s \\
&+ 2CL(t_f - s_0)\varrho\epsilon
\end{aligned}$$

Using Lemma 3.4, we have

$$g^*(t) \leq (\delta + 2CL(t_f - s_0)\varrho)\epsilon \prod_{s_i < s < t} \left(1 + \frac{2Lg_i}{m\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \Delta s \right) e_q(t, s_i).$$

where $q = 2CL \int_{s_0}^s \Delta u$. If we determine $g = \|\phi - \omega\|$, then $g(t) \leq (Tg)(t)$, which follows by utilizing abstract Grönwall lemma that $g(t) \leq g^*$, hence

$$\|\phi(t) - \omega(t)\| \leq (\delta + 2CL(t_f - s_0)\varrho)\epsilon \prod_{s_i < s < t} \left(1 + \frac{2Lg_i}{m\Gamma(\alpha)} \int_{t_i}^{s_i} (s_i - s)^{\alpha-1} \Delta s \right) e_q(t, s_i).$$

□

Similarly we can establish the Hyers–Ulam–Rassias stability of (1.1) on J . Its proof will be omitted.

Theorem 4.3. *If conditions $(\mathbf{A}_1) - (\mathbf{A}_4)$ hold, then Eq. (1.1) has Hyers–Ulam–Rassias stability on J .*

5. CONCLUSION

This manuscript is about the establishment of Hyers–Ulam stability and Hyers–Ulam–Rassias stability of equation (1.1) with the utilization of fixed point approach. Also, the unique solution to (1.1) in $PC^1(J, \mathbb{R}^n)$ is obtained. Furthermore, abstract Grönwall lemma and Lemma 3.4 presented a fruitful outcome to our end. Our work assures the existence of an exact solution of (1.1) near to approximate solution. In fact, our results are significant when finding exact solution is quite difficult and hence are important to approximation theory etc.

REFERENCES

- [1] R. P. Agarwal, A. S. Awan, D. ÓRegan and A. Younus, *Linear impulsive Volterra integro-dynamic system on time scales*, Adv. Difference Equ. **2014** (2014): 6
- [2] C. Alsina and R. Ger, *On some inequalities and stability results related to the exponential function*, J. Inequal. Appl. **2** (1998), 373–380.
- [3] S. András and A. R. Mészáros, *Ulam–Hyers stability of dynamic equations on time scales via Picard operators*, Appl. Math. Comput. **209** (2013), 4853–4864.
- [4] D. D. Bainov and A. Dishliev, *Population dynamics control in regard to minimizing the time necessary for the regeneration of a biomass taken away from the population*, Comp. Rend. Bulg. Sci. **42** (1989), 29–32.
- [5] D. D. Bainov and P. S. Simenov, *Systems with Impulse Effect Stability Theory and Applications*, Ellis Horwood Limited, Chichester, UK, 1989.
- [6] M. Bohner and A. Peterson, *Dynamic Equations on Time Scales: An Introduction with Applications*, Birkhäuser, Boston, Mass, USA, 2001.
- [7] M. Bohner and A. Peterson, *Advances in Dynamics Equations on Time Scales*, Birkhäuser, Boston, Mass, USA, 2003.
- [8] J. J. Dachunha, *Stability for time varying linear dynamic systems on time scales*, J. Comput. Appl. Math. **176** (2005), 381–410.
- [9] A. Hamza and K. M. Oraby, *Stability of abstract dynamic equations on time scales*, Adv. Difference Equ. **2012** (2012): 143.
- [10] S. Hilger, *Analysis on measure chains—A unified approach to continuous and discrete calculus*, Result math. **18** (1990), 18–56.
- [11] D. H. Hyers, *On the stability of the linear functional equation*, Proc. Nat. Acad. Sci. U.S.A. **27** (1941), 222–224.
- [12] S.-M. Jung, *Hyers–Ulam stability of linear differential equations of first order*, Appl. Math. Lett. **17** (2004), 1135–1140.
- [13] S.-M. Jung, *Hyers–Ulam–Rassias Stability of Functional Equations in Nonlinear Analysis*, Springer Optim. Appl., vol. 48, Springer, NewYork, 2011.
- [14] Y. Li and Y. Shen, *Hyers–Ulam stability of linear differential equations of second order*, Appl. Math. Lett. **23** (2010), 306–309.
- [15] Z. Lin, W. Wei and J. Wang, *Existence and stability results for impulsive integro-differential equations*, Ser. Math. Inform. **29** (2014), 119–130.
- [16] V. Lupulescu and A. Zada, *Linear impulsive dynamic systems on time scales*, Electron. J. Qual. Theory Differ. Equ. (2010), 1–30.
- [17] S. I. Nenov, *Impulsive controllability and optimization problems in population dynamics*, Nonlinear Anal. Theory Methods Appl. **36** (1999), 881–890.
- [18] M. Obloza, *Hyers stability of the linear differential equation*, Rocznik Nauk.-Dydakt. Prace Mat. (1993), 259–270.

- [19] M. Obłozza, *Connections between Hyers and Lyapunov stability of the ordinary differential equations*, Rocznik Nauk.-Dydakt. Prace Mat. **14** (1997), 141–146.
- [20] C. Pötzsche, S. Siegmund and F. Wirth, *A spectral characterization of exponential stability for linear time-invariant systems on time scales*, Discrete Contin. Dyn. Sys. **9** (2003), 1223–1241.
- [21] T. M. Rassias, *On the stability of linear mappings in Banach spaces*, Proc. Amer. Math. Soc. **72** (1978), 297–300.
- [22] R. Rizwan, J. R. Lee, C. Park and A. Zada, *Switched coupled system of nonlinear impulsive Langevin equations with mixed derivatives*, AIMS Math. **6** (2021), 13092–13118.
- [23] R. Rizwan and A. Zada, *Nonlinear impulsive Langevin equation with mixed derivatives*, Math. Methods Appl. Sci. **43** (2020), 427–442.
- [24] R. Rizwan and A. Zada, *Existence theory and Ulam’s stabilities of fractional Langevin equation*, Qual. Theory Dyn. Syst. **20** (2021), 1–17.
- [25] R. Rizwan, A. Zada, M. Ahmad, S. O. Shah and H. Waheed, *Existence theory and stability analysis of switched coupled system of nonlinear implicit impulsive Langevin equations with mixed derivatives*, Math. Methods Appl. Sci. **44** (2021), 8963–8985.
- [26] R. Rizwan, A. Zada and X. Wang, *Stability analysis of non linear implicit fractional Langevin equation with non-instantaneous impulses*, Adv. Differ. Equ. **2019** (2019), 1–31.
- [27] I. A. Rus, *Grönwall lemmas: ten open problems*, Sci. Math. Jpn. **70** (2009), 221–228.
- [28] S. O. Shah and A. Zada, *On the stability analysis of non-linear Hammerstein impulsive integro-dynamic system on time scales with delay*, Punjab Univ. J. Math. **51** (2019), 89–98.
- [29] S. O. Shah and A. Zada, *Existence, uniqueness and stability of solution to mixed integral dynamic systems with instantaneous and noninstantaneous impulses on time scales*, Appl. Math. Comput. **359**(2019), 202–213.
- [30] S. O. Shah and A. Zada, *Hyers–Ulam stability of non-linear Volterra integro-delay dynamic system with fractional integrable impulses on time scales*, Iran. J. Math. Sci., <http://ijmsi.ir/browse-accepted.php>(in press).
- [31] S. O. Shah, A. Zada and A. E. Hamza, *Stability analysis of the first order non-linear impulsive time varying delay dynamic system on time scales*, Qual. Theory Dyn. Syst. **18** (2019), 825–840.
- [32] S. O. Shah, A. Zada, M. Muzamil, M. Tayyab and R. Rizwan, *On the Bielecki–Ulam’s type stability results of first order non-linear impulsive delay dynamic systems on time scales*, Qual. Theory Dyn. Syst. **19** (2020), 1–18.
- [33] S. O. Shah, A. Zada, C. Tunc and A. Ali, *Bielecki–Ulam–Hyers stability of non-linear Volterra impulsive integro-delay dynamic systems on time scales*, Punjab Univ. J. Math. **53** (2021), 339–349.
- [34] S. M. Ulam, *A Collection of the Mathematical Problems*, Interscience Publisheres, New York-London, 1960.
- [35] S. M. Ulam, *Problem in Modern Mathematics*, Science Editions, J. Wiley and Sons, Inc., New York, 1964.
- [36] J. Wang, M. Feckan and Y. Zhou, *Ulam’s type stability of impulsive ordinary differential equations*, J. Math. Anal. Appl. **395** (2012), 258–264.
- [37] J. Wang, M. Feckan and Y. Zhou, *On the stability of first order impulsive evolution equations*, Opuscula Math. **34** (2014), 639–657.
- [38] X. Wang, R. Rizwan, J. R. Lee, A. Zada and S. O. Shah, *Existence, uniqueness and Ulam’s stabilities for a class of implicit impulsive Langevin equation with Hilfer fractional derivatives*, AIMS Math. **6**(2021), 4915–4929.
- [39] J. Wang and Y. Zhang, *A class of nonlinear differential equations with fractional integrable impulses*, Com. Nonl. Sci. Num. Sim. **19** (2014), 3001–3010.
- [40] J. Wang, A. Zada and W. Ali, *Ulam’s-type stability of first-order impulsive differential equations with variable delay in quasi-Banach spaces*, Int. J. Nonlin. Sci. Num. **19** (2018), 553–560.
- [41] A. Younus, D. O’Regan, N. Yasmin and S. Mirza, *Stability criteria for nonlinear volterra integro-dynamic systems*, Appl. Math. Inf. Sci. **11** (2017), 1509–1517.
- [42] A. Zada, R. Rizwan, J. Xu and Z. Fu, *On implicit impulsive Langevin equation involving mixed order derivatives*, Adv. Differ. Equ., **2019** (2019), 1–26.

- [43] A. Zada, W. Ali and S. Farina, *Hyers–Ulam stability of nonlinear differential equations with fractional integrable impulses*, Math. Methods Appl. Sci. **40** (2017), 5502–5514.
- [44] A. Zada, T. Li, S. Ismail and O. Shah, *Exponential dichotomy of linear autonomous systems over time scales*, Differ. Equ. Appl. **8** (2016), 123–134.
- [45] A. Zada, B. Pervaiz, S. O. Shah and J. Xu, *Stability analysis of first-order impulsive nonautonomous system on timescales*, Math. Methods Appl. Sci. **43** (2020), 5097–5113.
- [46] A. Zada and S. O. Shah, *Hyers–Ulam stability of first-order non-linear delay differential equations with fractional integrable impulses*, Hacettepe J. Math. Stat. **47** (2018), 1196–1205.
- [47] A. Zada, S. O. Shah, S. Ismail and T. Li, *Hyers–Ulam stability in terms of dichotomy of first order linear dynamic systems*, Punjab Univ. J. Math. **49** (2017), 37–47.
- [48] A. Zada, S. O. Shah and Y. Li, *Hyers–Ulam stability of nonlinear impulsive Volterra integro-delay dynamic system on time scales*, J. Nonlinear Sci. Appl. **10** (2017), 5701–5711.
- [49] A. Zada, O. Shah and R. Shah, *Hyers–Ulam stability of non-autonomous systems in terms of boundedness of Cauchy problems*, Appl. Math. Comput. **271** (2015), 512–518.

Manuscript received August 3 2021

revised October 29 2021

S. O. SHAH

Department of Physical and Numerical Sciences, Qurtuba University of Science and Information Technology, Peshawar 25000, Pakistan

E-mail address: omarshah89@yahoo.com; omarshah@qurtuba.edu.pk

Z. KHAN

Department of Physical and Numerical Sciences, Qurtuba University of Science and Information Technology, Peshawar 25000, Pakistan

E-mail address: zk6095180@gmail.com